

MATH553. Topology and Geometry of Surfaces
 Problem Sheet 2: Quotient Topology

Please hand in your solutions in class on *Thursday 13th October*. Question 3 is part of the assessment on this module. Office hours for this module are now fixed as: Mondays at 3, Tuesdays at 4, Fridays at 11, all in 515, which is reached through 516.

1. Let $X = \mathbb{R}^2 \setminus \{0\}$, and define \sim by: $\underline{x} \sim \underline{x}' \Leftrightarrow \underline{x}' = \lambda \underline{x}$ for some $\lambda > 0$. Check that \sim is an equivalence relation on X . Take the usual (subspace) topology on X , and the quotient topology on X/\sim . Take the usual (subspace) topology on $S^1 = \{(x, y) : x^2 + y^2 = 1\}$. Find a continuous map $F : X \rightarrow S^1$ such that $[F] : X/\sim \rightarrow S^1$ is well-defined and a bijection. (It will then automatically be continuous.)

2. Let $X = \mathbb{C} \times \{1, 2\}$. Give X the usual topology (as a subspace of \mathbb{C}^2) and let $\mathbb{C} \cup \{\infty\}$ be given the 1-point-compactification topology. Let $F : \mathbb{C} \times \{1, 2\} \rightarrow \mathbb{C} \cup \{\infty\}$ be defined by

$$F(z, 1) = z, \quad F(1/z, 2) = z \text{ if } z \neq 0, \quad F(0, 2) = \infty.$$

Show that F is continuous.

Hint: all you really need to show is that if $U \subset \mathbb{C}$ is an open set such that $\mathbb{C} \setminus U$ is bounded, then $\{1/z : z \in U\} \cup \{0\}$ is open in \mathbb{C} .

Now let the equivalence relation \sim be defined on X by: $(z, j) \sim (z', k) \Leftrightarrow$ either $(z, j) = (z', k)$ or $z' = 1/z$ and $j \neq k$. Check that \sim is an equivalence relation, and show that $[F] : X/\sim \rightarrow \mathbb{C} \cup \{\infty\}$ is well-defined, and a bijection. (It is then automatically continuous.)

3. *This problem is part of the CA components of this module and is worth 3 marks.*

For the letter or number you have been given, find a subset S_1 of \mathbb{R}^2 which, with the subspace topology, looks like the letter or number. Let

$$S_2 = \cup_{j=1}^n [a_j, b_j] \times \{j\} \subset \mathbb{R}^2$$

for some positive integer n , and intervals $[a_j, b_j]$ which you can choose at your convenience. Let S_2 be given the subspace topology, with respect to the standard topology on \mathbb{R}^2 . Choose an equivalence relation \sim such on S_2 that the quotient space S_2/\sim , with the quotient topology, is homeomorphic to S_1 , and find a homeomorphism $G : S_2/\sim \rightarrow S_1$, proving that it is indeed a homeomorphism.


Hint for last part Since S_2 is compact and S_1 is Hausdorff, it suffices to find a map $F : S_2 \rightarrow S_1$ which is continuous onto and such that $F(x_1) = F(x_2) \Leftrightarrow x_1 \sim x_2$.

Here is an example of how to tackle this problem for the letter X . There is more than one way to do this, and in fact the sets S_2 and S_2/\sim given below are not the same choices as made in lectures for the symbol $+$ — which is essentially the same as X .

Let

$$S_1 = \{(t, t) \in \mathbb{R}^2 : -1 \leq t \leq 1\} \cup \{(t, -t) \in \mathbb{R}^2 : -1 \leq t \leq 1\}.$$

This is a union of two line segments, one from $(-1, -1)$ to $(1, 1)$ which lies on the line $x = y$ and the other from $(-1, 1)$ to $(1, -1)$ which lies on the line $x + y = 0$. The two line segments intersect at $(0, 0)$. This certainly looks like the letter X .



Now let

$$S_2 = [-1, 1] \times \{1, 2\}.$$

Define \sim by $(0, 1) \sim (0, 2)$, and all other equivalence classes are trivial. Then we claim that S_2/\sim , with the quotient topology, is homeomorphic to S_1 . Since $S_1 \subset \mathbb{R}^2$, S_1 is Hausdorff, and since S_2 is a closed bounded subset of \mathbb{R}^2 , S_2 is compact, and the quotient S_2/\sim is compact, because the map $x \mapsto [x] : S_2 \rightarrow S_2/\sim$ is continuous onto, where $[x]$ is the equivalence class of x with respect to \sim . A continuous map from a compact space to a Hausdorff space which is also a bijection is a homeomorphism. So it suffices to find a continuous bijection $G : S_2/\sim \rightarrow S_1$. A continuous surjection $F : S_2 \rightarrow S_1$ such that $F^{-1}([x]) = \{y + y \sim x\} (= [x])$ gives rise to a continuous bijection $[F] : S_2/\sim \rightarrow S_1$ defined by $[F]([x]) = [F(x)]$, as shown in lectures. So it suffices to find such a continuous surjection F . We define F by

$$F(t, 1) = (t, t), F(t, 2) = (t, -t) \text{ for all } t \in [-1, 1].$$

Then $F : S_2 \rightarrow S_1$ is a surjection. Clearly $F(t, 1) = F(s, 1) \Leftrightarrow s = t$ and $F(t, 2) = F(s, 2) \Leftrightarrow s = t$. Also $F(t, 1) = F(s, 2) \Leftrightarrow (t, t) = (s, -s) \Leftrightarrow s = t = 0$. So F has all the required properties for $[F]$ to be well-defined and a homeomorphism.

4. Let the equivalence relation \sim be defined on \mathbb{C} by $z \sim z' \Leftrightarrow z' = z + m + ni$ for some $m, n \in \mathbb{Z}$. Check that this is an equivalence relation. Fix $\lambda \in \mathbb{C}$ and let $F : \mathbb{C} \rightarrow \mathbb{C}$ be given by $F(z) = \lambda z$. Show that $[F] : \mathbb{C}/\sim \rightarrow \mathbb{C}/\sim$ is well-defined $\Leftrightarrow \lambda = a + ib$ for some $a, b \in \mathbb{Z}$.

Also, determine for which λ $[F]$ is injective.

5. Let \sim be as in question 3. Let \approx be the equivalence relation defined on $\mathbb{C} \setminus \{0\}$ by $z' \approx z \Leftrightarrow z' = e^{2\pi n} z$ for some $n \in \mathbb{Z}$. Find a continuous map $F : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C} \setminus \{0\}$ such that $z \sim z' \Leftrightarrow F(z) \approx F(z')$, where F is *not* a bijection but $[F] : \mathbb{C}/\sim \rightarrow \mathbb{C}/\approx$ is .